

Supersymmetric Dark Matter and Recent Experimental Constraints

A. B. Lahanas¹, D. V. Nanopoulos², and V. C. Spanos³

¹ University of Athens, Physics Department, Nuclear and Particle Physics Section, GR-15771 Athens, Greece

² Department of Physics, Texas A & M University, College Station, TX 77843-4242, USA, Astroparticle Physics Group, Houston Advanced Research Center (HARC), Mitchell Campus, Woodlands, TX 77381, USA, and
Academy of Athens, Chair of Theoretical Physics, Division of Natural Sciences, 28 Panepistimiou Avenue, Athens 10679, Greece

³ Institut für Hochenergiephysik der Österreichischen Akademie der Wissenschaften, A-1050 Vienna, Austria

Abstract. In this talk we discuss the impact of recent experimental information, like the revised bound from E821 Brookhaven experiment on $g_\mu - 2$ and light Higgs boson mass bound from LEP, in delineating regions of the parameters of the Constrained Minimal Supersymmetric Standard Model which are consistent with the cosmological data. The effect of these to Dark Matter direct searches is also discussed.

ACT-03/02, CTP-TAMU-10/02

HEPHY-PUB 758/02

hep-ph/0205225

1 Introduction

Supersymmetry (SUSY) is not only indispensable in constructing consistent string theories, but it also seems unavoidable at low energies (~ 1 TeV) if the gauge hierarchy problem is to be resolved. Such a resolution provides a measure of the supersymmetry breaking scale $M_{SUSY} \approx \mathcal{O}(1 \text{ TeV})$. There is indirect evidence for such a low-energy supersymmetry breaking scale, from the unification of the gauge couplings [1] and from the apparent lightness of the Higgs boson as determined from precise electroweak measurements, mainly at LEP [2]. Furthermore, such a low energy SUSY breaking scale is also favored cosmologically, provided that R -parity is conserved. The spectrum of R -conserved SUSY models contain a stable, neutral particle, identifiable with the lightest neutralino ($\tilde{\chi}$), referred to as the LSP [3]. Such a particle is an ideal candidate for the Dark Matter (DM) in the Universe. The latest data from the Cosmic Microwave Background (CMB) radiation anisotropies [4] not only favour a flat ($k = 0$ or $\Omega_0 = 1$), inflationary Universe, but they also determine a matter density $\Omega_M h_0^2 \approx 0.15 \pm 0.05$. Subtracting from this the baryon density $\Omega_B h_0^2 \approx 0.02$, and the rather tiny neutrino density, one gets $\Omega_{DM} h_0^2 = 0.13 \pm 0.05$ for the DM density.

Assuming that all DM is supersymmetric due to LSP, i.e. $\Omega_{DM} \equiv \Omega_{\tilde{\chi}}$, one can combine the cosmological bound with other presently available constraints

from particle physics, such as the lower bound on the mass of the Higgs bosons ($m_h \geq 113.5$ GeV) provided by LEP [5] and the recent results from the BNL E821 experiment [6] on the anomalous magnetic moment of the muon ($\delta\alpha_\mu = 26(16) \times 10^{-10}$). The latter is the updated bound after the correction of a sign error in the hadronic light-by-light scattering contribution [7] to $g_\mu - 2$.

We find that the updated $g_\mu - 2$ bound does not put severe constraints on the parameter space of the Constrained Minimal Supersymmetric Standard Model (CMSSM) [8], especially in the 1.5σ region of this bound, in contrast to the previous situation [9, 10]. Nevertheless this new situation does not seem to affect significantly the potential of discovering SUSY at future direct DM searches.

2 Neutralino relic density

In the large $\tan\beta$ regime the neutralino ($\tilde{\chi}$) pair annihilation through s -channel pseudo-scalar Higgs boson (A) exchange, leads to an enhanced annihilation cross sections reducing significantly the relic density [11]. The importance of this mechanism, in conjunction with the recent cosmological data which favour small values of the DM relic density, has been stressed in [12, 13]. The same mechanism has been also invoked [14] where it has been shown that it enlarges the cosmologically allowed regions. In fact cosmology does not put severe upper bounds on sparticle masses, and soft masses can be in the TeV region, pushing up the sparticle mass spectrum to regions that might escape detection in future planned accelerators. Such an upper bound is imposed, however, by the $(g_\mu - 2)$ E821 data and has been the subject of intense phenomenological study the last year [9, 10, 15–18].

As was mentioned above, for large $\tan\beta$ the $\tilde{\chi}\tilde{\chi} \xrightarrow{A} b\bar{b}$ or $\tau\bar{\tau}$ channel becomes the dominant annihilation mechanism. In fact by increasing $\tan\beta$ the mass m_A decreases, while the neutralino mass remains almost constant, if the other parameters are kept fixed. Thus m_A is expected eventually to enter into the regime in which it is close to the pole value $m_A = 2m_{\tilde{\chi}}$, and the pseudo-scalar Higgs exchange dominates. It is interesting to point out that in a previous analysis of the direct DM searches [13], we had stressed that the contribution of the CP -even Higgs bosons exchange to the LSP-nucleon scattering cross sections increases with $\tan\beta$. Therefore in the large $\tan\beta$ region one obtains the highest possible rates for the direct DM searches. Similar results are presented in Ref. [19].

For the correct calculation of the neutralino relic density in the large $\tan\beta$ region, an unambiguous and reliable determination of the A -mass is required. The details of the procedure in calculating the spectrum of the CMSSM can be found elsewhere [9, 10]. Here we shall only briefly refer to some subtleties which turn out to be essential for a correct determination of m_A . In the CMSSM, m_A is not a free parameter but is determined once the other parameters are given. m_A depends sensitively on the Higgs mixing parameter, m_3^2 , which is determined from minimizing the one-loop corrected effective potential. For large $\tan\beta$ the derivatives of the effective potential with respect the Higgs fields, which enter into the minimization conditions, are plagued by terms which are large and hence potentially dangerous, making the perturbative treatment untrustworthy. In or-




Fig. 1. Cosmologically allowed regions of the relic density for two different values of $\tan\beta$ in the $(M_{1/2}, m_0)$ plane. The remaining inputs are shown in each figure. The mass of the top is taken 175 GeV. In the dark green shaded area $0.08 < \Omega_{\tilde{\chi}} h_0^2 < 0.18$. In the light green shaded area $0.18 < \Omega_{\tilde{\chi}} h_0^2 < 0.30$. The solid red lines mark the region within which the supersymmetric contribution to the anomalous magnetic moment of the muon is $\alpha_{\mu}^{SUSY} = (26 \pm 16) \times 10^{-10}$. The dashed red line is the boundary of the region for which the lower bound is moved to 1.5σ limit. The dashed-dotted blue lines are the boundaries of the region $113.5 \text{ GeV} \leq m_{Higgs} \leq 117.0 \text{ GeV}$.

der to minimize the large $\tan\beta$ corrections we had better calculate the effective potential using as reference scale the average stop scale $Q_{\tilde{t}} \simeq \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$ [20]. At this scale these terms are small and hence perturbatively valid. Also for the calculation of the pseudo-scalar Higgs boson mass all the one-loop corrections must be taken into account. In particular, the inclusion of those of the neutralinos and charginos yields a result for m_A that is scale independent and approximates the pole mass to better than 2% [21]. A more significant correction, which drastically affects the pseudo-scalar mass arises from the gluino-sbottom and chargino-stop corrections to the bottom quark Yukawa coupling h_b [22–25]. The proper resummation of these corrections is important for a correct determination of h_b [26,27], and accordingly of the m_A . In calculating the $\tilde{\chi}$ relic abundance, we solve the Boltzmann equation numerically using the machinery outlined in Ref. [12]. In this calculation the coannihilation effects, in regions where $\tilde{\tau}_R$ approaches in mass the LSP, which is a high purity Bino, are properly taken into account.

After the correction of the error in the sign of the hadronic light-by-light contribution to $g_{\mu} - 2$, actually there is no 2σ disagreement between the experimental value and the Standard Model prediction. There is still an 1σ discrepancy, and one can also study the 1.5σ bound. If we are going to attribute this discrepancy to SUSY the $\mu > 0$ case is relevant. In addition the $\mu < 0$ case is not favoured by the recent $b \rightarrow s\gamma$ data.

In the panels shown in figure 1 we display our results by drawing the cosmologically allowed region $0.08 < \Omega_{\tilde{\chi}} h_0^2 < 0.18$ (dark green) in the $m_0, M_{1/2}$

plane for values of $\tan\beta$ equal to 50 and 55 respectively. Also drawn (light green) is the region $0.18 < \Omega_{\tilde{\chi}} h_0^2 < 0.30$. In the figures shown we used for the top, tau and bottom masses the values $M_t = 175$ GeV, $M_\tau = 1.777$ GeV and $m_b(m_b) = 4.25$ GeV. The remaining inputs are shown on the top of each panel. The solid red mark the region within which the supersymmetric contribution to the anomalous magnetic moment of the muon falls within the E821 range $\alpha_\mu^{SUSY} = (26 \pm 16) \times 10^{-10}$. The dashed red line marks the boundary of the region when the more relaxed lower bound 1.5σ value of the E821 range. Along the blue dashed-dotted lines the light CP -even Higgs mass takes values 113.5 GeV (left) and 117.0 GeV (right) respectively. The line on the left marks therefore the recent LEP bound on the Higgs mass [5]. Also shown is the chargino mass bound 104 GeV. The shaded area (in red) at the bottom of each figure, labelled by TH, is theoretically disallowed since the light stau is lighter than the lightest of the neutralinos. From the displayed figures we observe that for values of $\tan\beta$ up to 50 the cosmological data put an upper bound on the parameter m_0 . However, there is practically no such upper bound for the parameter $M_{1/2}$, due to the coannihilation effects [14] which allow for $M_{1/2}$ as large as 1700 GeV within the narrow coannihilation band lying above the theoretically disallowed region.

For large values of $\tan\beta$, see the right panel of figure 1, a large region opens up within which the relic density is cosmologically allowed. This is due to the pair annihilation of the neutralinos through the pseudo-scalar Higgs exchange in the s -channel. As explained before, for such high $\tan\beta$ the ratio $m_A/2m_{\tilde{\chi}}$ approaches unity and the pseudo-scalar exchange dominates yielding large cross sections and hence small neutralino relic densities. In this case the lower bound put by the $g_\mu - 2$ data cuts the cosmologically allowed region which would otherwise allow for very large values of $m_0, M_{1/2}$.

For the $\tan\beta = 55$ case, close the highest possible value, and considering the lower bound on the muon's anomalous magnetic moment $\alpha_\mu^{SUSY} \geq 10 \times 10^{-10}$ and values of $\Omega_{\tilde{\chi}} h_0^2$ in the range 0.13 ± 0.05 , we find that the point with the highest value of m_0 is (in GeV) at $(m_0, M_{1/2}) = (950, 300)$ and that with the highest value of $M_{1/2}$ is at $(m_0, M_{1/2}) = (600, 800)$. The latter marks the lower end of the line segment of the boundary $\alpha_\mu^{SUSY} = 10 \times 10^{-10}$ which amputates the cosmologically allowed stripe. For the case displayed in the right panel of the figure 1 the upper mass limits put on the LSP, and the lightest of the charginos, stops and the staus are $m_{\tilde{\chi}} < 287, m_{\tilde{\chi}^+} < 539, m_{\tilde{t}} < 1161, m_{\tilde{\tau}} < 621$ (in GeV). Allowing for $A_0 \neq 0$ values, the upper bounds put on $m_0, M_{1/2}$ increase a little and so do the aforementioned bounds on the sparticle masses. Thus it appears that the prospects of discovering CMSSM at a e^+e^- collider with center of mass energy $\sqrt{s} = 800$ GeV, are not guaranteed. However in the allowed regions the next to the lightest neutralino, $\tilde{\chi}'$, has a mass very close to the lightest of the charginos and hence the process $e^+e^- \rightarrow \tilde{\chi}\tilde{\chi}'$, with $\tilde{\chi}'$ subsequently decaying to $\tilde{\chi} + l^+l^-$ or $\tilde{\chi} + 2\text{jets}$, is kinematically allowed for such large $\tan\beta$, provided the energy is increased to at least $\sqrt{s} = 900$ GeV. It should be noted however that this channel proceeds via the t -channel exchange of a selectron and it is suppressed due to the heaviness of the exchanged sfermion.




Fig. 2. Scatter plot of the scalar neutralino-nucleon cross section versus $m_{\tilde{\chi}}$, from a random sample of 40,000 points. On the top of the figure the CDMS excluded region and the DAMA sensitivity region are illustrated. Diamonds ($\diamond$) are points within the E821 experimental region $\alpha_{\mu}^{\text{SUSY}} = (26 \pm 16) \times 10^{-10}$ and which are cosmologically acceptable $\Omega_{\tilde{\chi}} h_0^2 = 0.13 \pm 0.05$. Crosses ($\times$) represent the rest of the random sample. The Higgs boson mass bound $m_h > 113.5$ GeV is properly taken into account.

The constraints put by the 1.5σ $g_{\mu} - 2$ bound are extremely weak, especially for large values of $\tan\beta$, as the $\alpha_{\mu}^{\text{SUSY}}$ is proportional to $\tan\beta$ [28]. Actually the 1.5σ region lies in the border line of LHC [29].

3 Direct Dark Matter searches

We turn now to study the impact of the $g_{\mu} - 2$ measurements and of the Higgs mass bound $m_h > 113.5$ GeV on the direct DM searches. For this reason we are using a random sample of 40,000 points in the region $|A_0| < 1$ TeV, $\tan\beta < 55$, $M_{1/2} < 1.5$ TeV, $m_0 < 1.5$ TeV and $\mu > 0$.

In figure 2 we plot the scalar $\tilde{\chi}$ -nucleon cross section as function of the LSP mass, $m_{\tilde{\chi}}$. On the top of the figure the shaded region (in cyan colour) is excluded by the CDMS experiment [30]. The DAMA sensitivity region (coloured in yellow) is also plotted [31]. Diamonds ($\diamond$) (in green colour) represent points which are both compatible with the E821 data $\alpha_{\mu}^{\text{SUSY}} = (26.0 \pm 16.0) \times 10^{-10}$ and the cosmological bounds for the neutralino relic density $\Omega_{\tilde{\chi}} h_0^2 = 0.13 \pm 0.05$. The crosses ($\times$) (in red colour) represent the rest of the points of our random sample. Here the Higgs boson mass bound, $m_h > 113.5$ GeV has been properly taken into account. From this figure it is seen that the points which are compatible both the $(g_{\mu} - 2)$ E821 and the cosmological data (crosses) yield cross sections of the order of 10^{-10} pb and the maximum value of the $m_{\tilde{\chi}}$ is about 350 GeV.

In figure 3 we don't impose the constraints stemming from $g_{\mu} - 2$ data, therefore due to the coannihilation processes the cosmologically acceptable LSP mass can be heavier than 500 GeV. It is worth noticing that that imposing the $g_{\mu} - 2$ bound the lowest allowed $\tilde{\chi}$ -nucleon cross section increases by about one




Fig. 3. Scatter plot of the scalar neutralino-nucleon cross section versus $m_{\tilde{\chi}}$, from a random sample of figure 2. Diamonds ($\diamond$) are cosmologically acceptable points, without putting any restriction from the $\alpha_{\mu}^{\text{SUSY}}$. Crosses ($\times$) represent points with unacceptable $\Omega_{\tilde{\chi}} h_0^2$.

order of magnitude, from 10^{-11} pb to 10^{-10} pb. Considering the $\mu > 0$ case, it is important that using the cosmological bound for the DM alone, one can put a lower bound on $\sigma_{\text{scalar}} \simeq 10^{-11}$. This fact is very encouraging for the future DM direct detection experiments [32]. Unfortunately this is not the case for $\mu < 0$, where the σ_{scalar} can be very small, due to an accidental cancellation between the sfermion and Higgs boson exchange processes. However, as discussed before, this case is not favoured by $b \rightarrow s\gamma$ data. Similar results are presented in Ref. [33].

4 Conclusions

In conclusion, we have combined recent high energy physics experimental information, like the anomalous magnetic moment of the muon measured at E821 Brookhaven experiment and the light Higgs boson mass bound from LEP, with the cosmological data for DM. Especially for the $g_{\mu} - 2$ bound we have used the revised value taking into account the correct sign in the hadronic light-by-light scattering contribution. We studied the imposed constraints on the parameter space of the CMSSM and hence we assessed the potential of discovering SUSY, if it is based on CMSSM, at future colliders and DM direct searches experiments. The use of the 1σ $g_{\mu} - 2$ bound can guarantee that in LHC but also in a e^+e^- linear collider with center of mass energy $\sqrt{s} = 1200$ GeV, CMSSM can be discovered. The 1.5σ bound is rather weak, resulting to a very heavy sparticle spectrum, especially for large $\tan\beta$. Actually this bound lies in the border line of LHC. The effect of these constraints is also significant for the direct DM searches. For the $\mu > 0$ case, even ignoring the $g_{\mu} - 2$ bound, we found that the

minimum value of the spin-independent $\tilde{\chi}$ -nucleon cross section attained is of the order of 10^{-11} pb.

Acknowledgements

A.B.L. acknowledges support from HPRN-CT-2000-00148 and HPRN-CT-2000-00149 programmes. He also thanks the University of Athens Research Committee for partially supporting this work. D.V.N. acknowledges support by D.O.E. grant DE-FG03-95-ER-40917. V.C.S. acknowledges support by a Marie Curie Fellowship of the EU programme IHP under contract HPMFCT-2000-00675. The authors thank the Academy of Athens for supporting this work.

References

1. J. Ellis, S. Kelley and D. V. Nanopoulos, Phys. Lett. B260 (1991) 131; U. Amaldi, W. de Boer and H. Furstenau, Phys. Lett. B260 (1991) 447; C. Giunti, C. W. Kim and U. W. Lee, Mod. Phys. Lett. A6 (1991) 1745.
2. LEP Electroweak Working Group, <http://lepewwg.web.cern.ch/LEPEWWG/Welcome.html>.
3. J. Ellis, J. S. Hagelin, D. V. Nanopoulos, K. A. Olive and M. Srednicki, Nucl. Phys. B238 (1984) 453; H. Goldberg, Phys. Rev. Lett. 50 (1983) 1419.
4. A. T. Lee *et al.*, MAXIMA Collaboration, Astrophys. J. 561 (2001) L1–L6, astro-ph/0104459; C. B. Netterfield *et al.*, BOOMERANG Collaboration, astro-ph/0104460; N. W. Halverson *et al.*, DASI Collaboration, astro-ph/0104489; P. de Bernardis *et al.*, Astrophys. J. 564 (2002) 559, astro-ph/0105296.
5. LEP Collaborations, CERN-EP/2001-055 (2001), hep-ex/0107021; LEP Higgs Working Group, hep-ex/0107030, <http://lephiggs.web.cern.ch/LEPHIGGS>.
6. H. N. Brown *et al.*, Muon $g_\mu - 2$ Collaboration, Phys. Rev. Lett. 86 (2001) 2227.
7. M. Knecht and A. Nyffeler, Phys. Rev. D65 (2002) 073034; M. Knecht, A. Nyffeler, M. Perrottet and E. de Rafael, Phys. Rev. Lett. 88 (2002) 071802; M. Hayakawa and T. Kinoshita, hep-ph/0112102.
8. J. Ellis, astro-ph/0204059; contributions to these proceedings by R. Arnowitt, J. Ellis, P. Nath and L. Roszkowski.
9. A. B. Lahanas and V. C. Spanos, Eur. Phys. J. C23 (2002) 185.
10. A. B. Lahanas, D. V. Nanopoulos and V. C. Spanos, Phys. Lett. B518 (2001) 94, hep-ph/0112134.
11. M. Drees and M. Nojiri, Phys. Rev. D47 (1993) 376; R. Arnowitt and P. Nath, Phys. Lett. B299 (1993) 58.
12. A. B. Lahanas, D. V. Nanopoulos and V. C. Spanos, Phys. Rev. D62 (2000) 023515.
13. A. B. Lahanas, D. V. Nanopoulos and V. C. Spanos, Mod. Phys. Lett. A16 (2001) 1229.
14. J. Ellis, T. Falk, G. Ganis, K. A. Olive and M. Srednicki, Phys. Lett. B510 (2001) 236.
15. J. Ellis, D. V. Nanopoulos and K. A. Olive, Phys. Lett. B508 (2001) 65;
16. L. Everett, G. L. Kane, S. Rigolin and L. Wang, Phys. Rev. Lett. 86 (2001) 3484; J. L. Feng and K. T. Matchev, Phys. Rev. Lett. 86 (2001) 3480; E. A. Baltz and P. Gondolo, Phys. Rev. Lett. 86 (2001) 5004; U. Chattopadhyay and P. Nath, Phys. Rev. Lett. 86 (2001) 5854; S. Komine, T. Moroi and M. Yamaguchi, Phys. Lett.

- B506 (2001) 93; J. Hisano and K. Tobe, Phys. Lett. B510 (2001) 197; R. Arnowitt, B. Dutta, B. Hu and Y. Santoso, Phys. Lett. B505 (2001) 177; K. Choi, K. Hwang, S. K. Kang, K. Y. Lee and W. Y. Song, Phys. Rev. D64 (2001) 055001; S. P. Martin and J. D. Wells, Phys. Rev. D64 (2001) 035003; S. Komine, T. Moroi and M. Yamaguchi, Phys. Lett. B507 (2001) 224; H. Baer, C. Balazs, J. Ferrandis and X. Tata, Phys. Rev. D64 (2001) 035004; V. Barger and C. Kao, Phys. Lett. B518 (2001) 117.
17. L. Roszkowski, R. R. Austri and T. Nihei, JHEP (2001) 0108:024.
 18. A. Djouadi, M. Drees and J. L. Kneur, JHEP (2001) 0108:055.
 19. M. Drees, Y. G. Kim, T. Kobayashi and M. M. Nojiri, Phys. Rev. D63 (2001) 115009.
 20. B. de Carlos and J. A. Casas, Phys. Lett. B309 (1993) 320; H. Baer, C. Chen, M. Drees, F. Paige and X. Tata, Phys. Rev. Lett. 76 (1997) 986; J. A. Casas, J. R. Espinosa and H. E. Haber, Nucl. Phys. B526 (1998) 3.
 21. A. Katsikatsou, A. B. Lahanas, D. V. Nanopoulos and V. C. Spanos, Phys. Lett. B501 (2001) 69.
 22. R. Hempfling, Phys. Rev. D49 (1994) 6168; L. J. Hall, R. Rattazi and U. Sarid, Phys. Rev. D50 (1994) 7048; R. Rattazi and U. Sarid, Phys. Rev. D53 (1996) 1553.
 23. M. Carena, M. Olechowski, S. Pokorski and C. Wagner, Nucl. Phys. B426 (1994) 269.
 24. J. A. Bagger, K. Matchev, D. M. Pierce and R. -J. Zhang, Nucl. Phys. B491 (1997) 3.
 25. E. Accomando, R. Arnowitt, B. Dutta and Y. Santoso, Nucl. Phys. B585 (2000) 124.
 26. H. Eberl, K. Hidaka, S. Kraml, W. Majerotto and Y. Yamada, Phys. Rev. D62 (2000) 055006.
 27. M. Carena, D. Garcia, U. Nierste and C. Wagner, Nucl. Phys. B577 (2000) 88.
 28. J. L. Lopez, D. V. Nanopoulos and X. Weng, Phys. Rev. D49 (1994) 366.
 29. U. Chattopadhyay and P. Nath, hep-ph/0101001.
 30. R. Abusaidi *et al.*, CDMS Collaboration, Phys. Rev. Lett. 84 (2000) 5699.
 31. R. Bernabei *et al.*, DAMA Collaboration, Phys. Lett. B480 (2000) 23.
 32. H. V. Klapdor-Kleingrothaus, hep-ph/0104028.
 33. R. Arnowitt, contribution to these proceedings, hep-ph/0204187; J. Ellis, A. Ferstl and K. A. Olive, hep-ph/0111064.